

HESSIAN OF THE ZETA FUNCTION FOR THE LAPLACIAN ON FORMS

K. OKIKIOLU AND C. WANG

ABSTRACT. Let M be a compact closed n -dimensional manifold. Given a Riemannian metric on M , we consider the zeta functions $Z(s)$ for the de Rham Laplacian and the Bochner Laplacian on p -forms. The hessian of $Z(s)$ with respect to variations of the metric is given by a pseudodifferential operator T_s . When the real part of s is less than $n/2 - 1$ we compute the principal symbol of T_s . This can be used to determine whether a general critical metric for $(d/ds)^k Z(s)$ has finite index, or whether it is an essential saddle point.

1. Introduction.

Over the last fifteen years, several results have been proved which identify maximal or minimal metrics for the determinant of a geometric elliptic operator on a compact manifold. In [OPS1], surfaces of constant curvature were shown to maximize the determinant of the Laplacian within the conformal class. Most higher dimensional cases studied involve extremizing a conformally covariant operator within a conformal class. In these cases local variation formulas are available, [BØ], [Po], and global results have been obtained, see [BCY], [Br], [CY]. (Some results also exist for manifolds which are non-closed or non-compact, see for example [BFKM], [CQ], [HZ], [OPS2], [OPS3].) In addition, extremal metrics for special values of the zeta function for conformally covariant operators have been studied, see [Mo]. The zeta function for the de Rham Laplacian on forms is interesting because of its connection to topology. For the Laplace-Beltrami operator Δ_0 , the standard 3-sphere is a local maximum for $\det \Delta_0$, but if $n \geq 7$ is odd, the standard n -sphere is a saddle for $\det \Delta_0$, see [Ok1], [Ri]. Beyond the scalar case, the determinant of the de Rham Laplacian was studied in relation to analytic torsion, [Ch], [Mü], [RS] and the variation of the zeta function was studied in [Ro], but there does not seem to have been progress on the subject of critical metrics since then. (In the non-scalar case there is not a great deal known about the spectrum of the Laplacian on forms. See [Ba], [BGV], [CT], [CC], [Fe], [Do], [Go], [Lo], [Ta] for results and further references.) In [Ok2] it was shown that for general geometric operators of Laplace type, the Hessian of the zeta function is given by a pseudodifferential operator T_s , and when $\Re s < n/2 - 1$, a formula for the principal symbol $u_s(x, \xi)$ was given. In the current paper we compute u_s explicitly for the case of the de Rham and Bochner Laplacians on p -forms. This can be applied to compute the behavior of the determinant and zeta function in the neighborhood of critical metrics. The computation of u_s is somewhat lengthy for the de Rham Laplacian. The proof given here can serve as a blueprint for carrying out the calculation for any operator of Laplace type. Happily, the results of the calculations are simple to state, see Theorem 1.

In order to present the results we set up some notation. Let M be a closed, compact n -dimensional manifold, let χ be a character of the fundamental group of M , let L_χ denote the complex line over M twisted by χ , let $\Omega^p(M, L_\chi)$ denote the space of smooth p -forms valued in L_χ , and let d denote the

The first author was supported by the National Science Foundation #DMS-9703329.

exterior differential. For the Riemannian metric g on M , let dV be the volume form on M and let V be the volume of M . For any tensor bundle E over M with values in L_χ , the metric g naturally gives rise to an inner product $\langle \cdot, \cdot \rangle_g$ on E , an inner product $\langle\langle \cdot, \cdot \rangle\rangle_g$ on the smooth sections of E , and a connection ∇ on E . We consider the de Rham Laplacian $d^*d + dd^*$ and the Bochner Laplacian $\nabla^*\nabla$ on $\Omega^p(M, L_\chi)$. These differ by a curvature operator. We use the notation Δ_p to denote either of the two operators. Let

$$\lambda_1 \leq \lambda_2 \leq \dots$$

be the non-zero eigenvalues of Δ_p . The spectral zeta function

$$Z(s) = Z(\Delta_p)(s) = \sum_{j=1}^{\infty} \lambda_j^{-s}$$

converges for $\Re s > n/2$ and extends to a meromorphic function with simple poles located at

$$s = \begin{cases} n/2, n/2 - 1, \dots, 1, & n \text{ even,} \\ n/2, n/2 - 1, \dots, \dots, & n \text{ odd.} \end{cases}$$

The spectral determinant of Δ_p is defined to be

$$\det \Delta_p = e^{-Z'(0)}.$$

Define the modified zeta function to be

$$(1.1) \quad \mathcal{Z}(s) = \mathcal{Z}(\Delta_p)(s) = \frac{\Gamma(s)Z(s)}{\Gamma(s - n/2)} + \frac{\dim \ker \Delta_p}{s \Gamma(s - n/2)},$$

which is entire in s . We remark that the dimension, $\beta_p(\chi)$, of the kernel of the de Rham Laplacian is a topological invariant, and is consequently invariant under changes of the metric.

Let S^2M denote the bundle of symmetric tensor fields of type $(0, 2)$ on M , and let $C^\infty(S^2M)$ denote its space of smooth sections. The Hessian of the modified zeta function is the bilinear form on $C^\infty(S^2M)$ defined by

$$\text{Hess } \mathcal{Z}(s)(h, h) = \left. \frac{d^2}{d\alpha^2} \right|_{\alpha=0} \mathcal{Z}(\Delta_p(g + \alpha h))(s).$$

We recall the following result which holds for the general geometric operator Δ of Laplace type defined on a tensor bundle.

[Ok2, Theorem 1]. *For $s \in \mathbb{C}$, there exists a unique symmetric pseudodifferential operator $T_s = T_s(\Delta) : C^\infty(S^2M) \rightarrow C^\infty(S^2M)$ such that*

$$(1.2) \quad \text{Hess } \mathcal{Z}(s)(h, h) = \langle\langle h, T_s h \rangle\rangle_g.$$

The operator T_s is analytic in s . For $s \notin n/2 + \mathbb{N}^+$, there exist polyhomogeneous pseudodifferential operators U_s and V_s of degrees $n - 2s$ and 2 respectively such that $T_s = U_s + V_s$. The operators U_s and V_s are meromorphic in s with simple poles located in $n/2 + \mathbb{N}^+$. (The poles of U_s and V_s cancel in the sum $T_s = U_s + V_s$, but the symbol expansion of T_s for $s \in n/2 + \mathbb{N}^+$ involves logarithmic terms.) For general s , the polyhomogeneous symbol expansion of U_s is computable from the complete symbol of the

operator Δ . In particular, there is a simple algorithm to compute the term u_s of homogeneity $n - 2s$. Furthermore, we can differentiate in s to obtain

$$(1.3) \quad \text{Hess}(d/ds)^k \mathcal{Z}(s)(h, h) = \langle\langle h, (d/ds)^k T_s h \rangle\rangle_g,$$

and the principal symbol of $(d/ds)^k U_s$ is equal to the leading order term of $(d/ds)^k u_s$ (provided this does not vanish identically).

The purpose of this paper is to compute the symbol $u_s(x, \xi)$ for the de Rham and Bochner Laplacians. This is the principal symbol of T_s when $\Re s < n/2 - 1$, (providing it does not vanish identically) and so for s in this range, u_s governs to some extent how $(d/ds)^k \mathcal{Z}(s)$ behaves in the neighborhood of a critical metric. In particular, $u_s(x, \xi)$ generally governs the behavior of $\log \det \Delta$. We remark that the above theorem was stated in [Ok2] for the case when the Laplacian has been normalized to be scale invariant, but this assumption is easily seen to be unnecessary.

For a point $x \in M$, we write $\text{End}(T_x^* M)$ for the endomorphisms of the cotangent space $T_x^* M$, and we write “tr” for the trace function on $\text{End}(T_x^* M)$. It is convenient to introduce the variable $S = s - n/2$, and set

$$(1.4) \quad C = C(n, s) = \left(\frac{1}{4\pi} \right)^{n/2} \frac{\Gamma(-S+1)^2}{\Gamma(-2S+2)}.$$

Theorem 1. *Suppose g is a metric on M , x is a point of M , and $h \in S_x^2 M$. Set*

$$(1.5) \quad H = hg^{-1} \in \text{End}(T_x^* M).$$

For a non-zero covector $\xi \in T_x^ M$, let*

$$\Pi_\xi : T_x^*(M) \rightarrow T_x^*(M)$$

denote the g -orthogonal projection onto the space spanned by ξ , and let

$$(1.6) \quad \Pi_\xi^\perp = I - \Pi_\xi.$$

Let Δ_p be the de Rham or Bochner Laplacian on p -forms, and in the case of the Bochner Laplacian assume that the dimension of its kernel has constant dimension when g is deformed. Then at the metric g ,

$$(1.7) \quad \langle h, u_s(x, \xi) h \rangle_g = 2C(n, s) |\xi|^{n-2s} \left(f_1(\Delta_p) \text{tr}((H \Pi_\xi^\perp)^2) + f_2(\Delta_p) (\text{tr} H \Pi_\xi^\perp)^2 \right),$$

where

$$(1.8) \quad f_1(\nabla^* \nabla)(n, p, s) = 4 \binom{n-2}{p-1} (S - 1/2) + \frac{1}{2} \binom{n}{p},$$

$$(1.9) \quad f_2(\nabla^* \nabla)(n, p, s) = \binom{n}{p} (S^2 + S - 3/4) + \frac{1}{4} \binom{n}{p}.$$

$$(1.10) \quad f_1(d^* d + dd^*)(n, p, s) = 4 \binom{n-2}{p-1} (S^2 + S - 3/4) + \frac{1}{2} \binom{n}{p},$$

$$(1.11) \quad f_2(d^* d + dd^*)(n, p, s) = \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) (S^2 + S - 3/4) + \frac{1}{4} \binom{n}{p},$$

Before discussing applications, we point out some features of the result. Firstly one notices that for $\xi, \eta \in T_x^*M$, if $h \in S_x^2M$ is given by

$$(1.12) \quad h = \xi \otimes \eta + \eta \otimes \xi$$

then

$$\Pi_\xi^\perp H \Pi_\xi^\perp = 0.$$

Hence for any $k \in S_x^2M$, writing $K = kg^{-1}$ we have

$$(1.13) \quad \langle k, u_s(x, \xi)h \rangle_g = 2C(n, s)|\xi|^{n-2s} \left(f_1 \text{trace} (K \Pi_\xi^\perp H \Pi_\xi^\perp) + f_2 \left(\text{trace} K \Pi_\xi^\perp \right) \left(\text{trace} H \Pi_\xi^\perp \right) \right) = 0.$$

This reflects on a micro-local level the fact that if $\phi(\alpha)$ is a one parameter family of diffeomorphisms of M with $\phi(0)$ equal to the identity and we set

$$h = \left. \frac{d\phi(\alpha)^*g}{d\alpha} \right|_{\alpha=0},$$

then $\text{Hess } \mathcal{Z}(s)(k, h)$ vanishes for every $k \in C^\infty(S^2M)$. It is easy to see that if $h \in S^2M$ does not have the form in (1.12) then

$$(1.14) \quad \text{trace} ((H \Pi_\xi^\perp)^2) > 0, \quad \frac{1}{n-1} \text{trace} ((H \Pi_\xi^\perp)^2) \geq \left(\text{trace} H \Pi_\xi^\perp \right)^2 \geq 0.$$

Secondly, we remark that instead of considering the full de Rham Laplacian, we can consider the operator d^*d acting on p -forms. The zeta function for this operator is

$$Z((d^*d)_p)(s) = \sum_{q=0}^p (-1)^{p-q} Z((d^*d + dd^*)_q)(s).$$

From Theorem 1, we find that the symbol $u_s(x, \xi)$ for the Hessian of $\mathcal{Z}_{d^*d}(s)$ is given by

$$(1.15) \quad \langle h, u_s(x, \xi)h \rangle_g = C(n, s) \left(f_1(n-1, p, s) \text{tr} ((H \Pi_\xi^\perp)^2) + f_2(n-1, p, s) \left(\text{tr} H \Pi_\xi^\perp \right)^2 \right) |\xi|^{n-2s}.$$

Thirdly we remark that the analytic torsion τ defined in the acyclic case by

$$\frac{1}{2} \log \tau = \sum_{p=0}^n (-1)^{p+1} p \log \det(d^*d + dd^*)_p$$

is a topological invariant. This implies that the corresponding alternating sum of symbols $u_0(x, \xi)$ for $d^*d + dd^*$ on p forms must vanish, and this is easily seen to be the case. In fact a stronger result is easily seen to be true from Theorem 1, namely for both the de Rham and the Bochner Laplacian Δ_p and for all s and all $k = 0, 1, \dots$,

$$(1.16) \quad \sum_{p=0}^n (-1)^{p+1} p^k u_s(\Delta_p)(x, \xi) = 0.$$

Lastly we remark that the character χ plays no role in our results which come from local calculations. The only thing one sees in the results related to the line bundle L_χ is the factor “2” in the right hand side of (1.7) which comes from the fact that the complex line has real dimension 2.

In a subsequent paper, we will show how Theorem 1 implies following results.

Theorem 2. Suppose $n \geq 3$, s is real and $s < n/2 - 1$, and Δ_p is either the de Rham or the Bochner Laplacian. For $j = 1, 2$, we consider the real numbers $f_j = f_j(\Delta_p)(n, p, s)$.

(a). If f_1 and $f_1/(n-1) + f_2$ are both positive or both negative, then every critical metric g for

$$(1.17) \quad (-1)^k \operatorname{sgn}(f_1) (d/ds)^k \mathcal{Z}(s)$$

has finite index. (That is, there exists a submanifold $\mathcal{X}$ of $\mathcal{M}$ of finite codimension, containing g , such $(-1)^k \operatorname{sgn}(f_1) (d/ds)^k \mathcal{Z}(s)$ has a local minimum at g on $\mathcal{X}$.)

(b). Conversely, if f_1 and $f_1/(n-1) + f_2$ have opposite signs, then every critical metric g for $(d/ds)^k \mathcal{Z}(s)$ is an essential saddle point. (That is, there are two infinite dimensional submanifolds $\mathcal{X}, \mathcal{Y}$, of $\mathcal{M}$, both containing g , such that $(d/ds)^k \mathcal{Z}(s)$ has a local minimum at g on $\mathcal{X}$ and a local maximum at g on $\mathcal{Y}$.)

Corollary. Let $n \geq 3$. If $n \equiv 1 \pmod{4}$, every critical metric for $\det(d^*d + dd^*)_p$ has finite index. If $n \equiv 3 \pmod{4}$, every critical metric for $1/\det(d^*d + dd^*)_p$ has finite index. The same is true if the de Rham Laplacian is replaced by the operator d^*d or the operator dd^* acting on p forms. However, for the Bochner Laplacian on forms of degree $p \neq 0, n$, the determinant may have an essential saddle point. The lowest dimension in which this happens is $n = 4$.

We remark that because we have not normalized the Laplacian to be scale invariant, a critical metric in the results above is one which is critical for the given functional under variations of the metric which preserve the total volume.

The rest of the paper is devoted to proving Theorem 1. We work throughout in the real case, where the Laplacian is acting on real valued p -forms. Because the calculation is entirely local, the character χ plays no role. The effect of going from the real case to the case of the complex line L_χ is just to multiply u_s by the factor 2, and this is the factor 2 which appears in (1.7). In Section 2, we discuss the general formula for u_s which was given in [Ok2], and we rewrite it in terms of the coefficient symbols of the first variation of the Laplacian. In Section 3, we compute the leading order terms in the first variation of Δ_p , where Δ_p is the Bochner or de Rham Laplacian. This is carried out in coordinates utilizing (3.7) as suggested by a referee for an early version of the paper which dealt only with the de Rham operator. Our change in notation has the advantage that the Bochner and de Rham Laplacians can be dealt with in the same way, and it will also hopefully make the proofs easier to follow. In Section 4, we identify the matrix entries of the first variation of the Laplacian. In Sections 5 and 6, we substitute these matrix entries into our formula for u_s , thus proving Theorem 1. For case of the de Rham Laplacian, it is essential to work in coordinates ξ in which h is diagonal, since otherwise the calculation becomes extremely lengthy. It is justified to choose to work in such coordinates because u_s is the principal symbol of a pseudodifferential operator, and hence does not depend on the choice of coordinates. Section 7 contains some observations on the calculation of u_s for the de Rham Laplacian in general coordinates.

2. The general formula for u_s .

Our starting point to prove Theorem 1, is the following result which gives u_s in the general case.

[Ok2, Theorem 2]. The symbol u_s for the general geometric operator Δ of Laplace type on the tensor bundle E of real dimension N can be computed as follows. Write $\Delta'_p = (d/d\sigma)|_{\sigma=0} \Delta_p(g + \sigma h)$ and let $x \in M$. Take coordinates on M which are orthonormal in the metric g at the point x , and take a local trivialization of E on a neighborhood of x to obtain coordinates for E . Suppose that in these coordinates, the operator Δ'_p is given at the point x by

$$(2.1) \quad \Delta'_p = \sum_{\alpha, \beta, i, j} A_{\alpha, \beta}^{ij} (\partial_w^\alpha h_{ij}(x)) \partial_w^\beta,$$

where α and β are multiindices and $A_{\alpha,\beta}^{ij}$ is an $N \times N$ real valued matrix. Set

$$(2.2) \quad C(n, s) = \left(\frac{1}{4\pi} \right)^{n/2} \frac{\Gamma(-S+1)^2}{\Gamma(-2S+2)}.$$

Then at $(x, \xi) \in T^*M$, the value of $u_s(x, \xi) \in \text{End}(S^2 M)_x$ is given by

$$(2.3) \quad (u_s(x, \xi)h)_{ij} = C(n, s) \sum_{\substack{|\alpha|+|\beta|=2 \\ |\gamma|+|\delta|=2 \\ k, \ell}} u_s(\partial^\alpha, \partial^\beta, \partial^\gamma, \partial^\delta, x, \xi) \text{trace}(A_{\alpha,\beta}^{ij}(x) A_{\gamma,\delta}^{k\ell}(x)) h_{k\ell},$$

where writing I for the identity operator on E_x , the terms $u_s(\partial^\alpha, \partial^\beta, \partial^\gamma, \partial^\delta, x, \xi)$ are given as follows:

$$(2.4) \quad u_s(\partial_j \partial_k, I, \partial_p \partial_q, I, x, \xi) = 4(4S^2 - 1) \xi_j \xi_k \xi_p \xi_q |\xi|^{n-2s-4}.$$

$$(2.5) \quad u_s(\partial_j, \partial_k, \partial_p \partial_q, I, x, \xi) = -2(4S^2 - 1) \xi_j \xi_k \xi_p \xi_q |\xi|^{n-2s-4}.$$

$$(2.6) \quad u_s(\partial_j, \partial_k, \partial_p, \partial_q, x, \xi) = (4S^2 + 2S - 2) \xi_j \xi_k \xi_p \xi_q |\xi|^{n-2s-4} - (2S - 1) \delta_{kq} \xi_j \xi_p |\xi|^{n-2s-2}.$$

$$(2.7) \quad u_s(I, \partial_j \partial_k, \partial_p \partial_q, I, x, \xi) = (4S^2 - 2S) \xi_j \xi_k \xi_p \xi_q |\xi|^{n-2s-4} + (2S - 1) \delta_{jk} \xi_p \xi_q |\xi|^{n-2s-2}.$$

$$(2.8) \quad u_s(I, \partial_j \partial_k, \partial_p, \partial_q, x, \xi) = -(2S^2 + S - 1) \xi_j \xi_k \xi_p \xi_q |\xi|^{n-2s-4} \\ + (S - 1/2)(-\delta_{jk} \xi_p \xi_q + \delta_{jq} \xi_k \xi_p + \delta_{kq} \xi_j \xi_p) |\xi|^{n-2s-2}$$

$$(2.9) \quad u_s(I, \partial_j \partial_k, I, \partial_p \partial_q, x, \xi) = \\ (S^2 + S) \xi_j \xi_k \xi_p \xi_q |\xi|^{n-2s-4} + \frac{1}{2}(S - 1) \delta_{jk} \xi_p \xi_q |\xi|^{n-2s-2} + \frac{1}{2}(S - 1) \xi_j \xi_k \delta_{pq} |\xi|^{n-2s-2} \\ - \frac{S}{2}(\delta_{jp} \xi_k \xi_q + \delta_{kq} \xi_j \xi_p + \delta_{jq} \xi_k \xi_p + \delta_{kp} \xi_j \xi_q) |\xi|^{n-2s-2} + \frac{1}{4}(\delta_{jk} \delta_{pq} + \delta_{jp} \delta_{kq} + \delta_{jq} \delta_{kp}) |\xi|^{n-2s}.$$

It is convenient to introduce the following notation.

$$(2.10) \quad h \cdot \xi = \sum_i h_{ij} \xi_j, \quad \xi \cdot h \cdot \xi = \sum_{i,j} h_{ij} \xi_i \xi_j, \quad |h| = \left(\sum_{i,j} h_{ij}^2 \right)^{1/2}, \quad \text{tr } h = \sum_i h_{ii}.$$

Note that the matrix for $H\Pi_\xi^\perp$ is

$$h_{ij} = |\xi|^{-2} \sum_k h_{ik} \xi_k \xi_j,$$

so

$$(2.11) \quad \begin{aligned} \text{tr} \left((H \Pi_\xi^\perp)^2 \right) &= |h|^2 - 2|\xi|^{-2} |h \cdot \xi|^2 + |\xi|^{-4} (\xi \cdot h \cdot \xi)^2, \\ (\text{tr} H \Pi_\xi^\perp)^2 &= |\xi|^{-4} (\xi \cdot h \cdot \xi)^2 - 2|\xi|^{-2} (\xi \cdot h \cdot \xi) (\text{tr} h) + (\text{tr} h)^2. \end{aligned}$$

We will now write u_s as a sum

$$(2.12) \quad u_s = C(n, s) \left(u_s^{(1)} + u_s^{(2)} + u_s^{(3)} + u_s^{(4)} \right).$$

To define the symbols $u_s^{(j)}$, choose coordinates which are orthonormal for the metric g at the point x . We notice that (2.3) only involves those terms in (2.1) which have *total degree* 2, where the *total degree* of a term

$$(2.13) \quad (\partial_w^\alpha h_{ij}(x)) \partial_w^\beta$$

is $|\alpha| + |\beta|$. We write

$$(2.14) \quad \Delta' \sim \sum_{i,j} h_{ij} \partial_{ij}^2 + \sum_{i,j,k,\ell} \left(A_{k\ell}^{ij} (\partial_k h_{ij}) \partial_\ell + B_{k\ell}^{ij} (\partial_{k\ell}^2 h_{ij}) \right),$$

where the notation “ $\sim$ ” means that the expressions are equal up to terms having lower total degree, and where for i, j, k, ℓ fixed, $A_{k\ell}^{ij}$ and $B_{k\ell}^{ij}$ are $N \times N$ matrix valued functions. The term $u^{(1)}$ is obtained by considering the terms in (2.4)–(2.9) of the form $(S^2 - 1/4) \xi_j \xi_k \xi_p \xi_q$. We define the *coefficient symbols* of Δ' by replacing each occurrence of ∂_j in (2.14) by ξ_j .

$$(2.15) \quad \sigma^{(2)} = \sum_{i,j} h_{ij} \xi_i \xi_j, \quad \sigma^{(1)} = \sum_{i,j,k,\ell} A_{k\ell}^{ij} \xi_k \xi_\ell h_{ij} \quad \sigma^{(0)} = \sum_{i,j,k,\ell} B_{k\ell}^{ij} \xi_k \xi_\ell h_{ij}.$$

These are $N \times N$ matrix valued. We also define the *coefficient components* of Δ' by

$$(2.16) \quad \sigma_{k\ell}^{(1)} = \sum_{i,j} A_{k\ell}^{ij} h_{ij} \quad \sigma_{k\ell}^{(0)} = \sum_{i,j} B_{k\ell}^{ij} h_{ij}.$$

Then $u_s^{(1)}$ is obtained by collecting all multiples of $(S^2 - 1/4) \xi_j \xi_k \xi_p \xi_q$ from (2.4)–(2.9), and its quadratic form is given by

$$(2.17) \quad \langle h, (u_s^{(1)}(x, \xi) h) \rangle := (S^2 - 1/4) |\xi|^{n-2s-4} \text{trace} \left(\left(\sigma^{(2)} - 2\sigma^{(1)} + 4\sigma^{(0)} \right)^2 \right).$$

The symbol $u_s^{(2)}$ is obtained from (2.6) after subtracting the term which goes into $u_s^{(1)}$. Its quadratic form is given by

$$(2.18) \quad \langle h, (u_s^{(2)}(x, \xi) h) \rangle = (2S - 1) |\xi|^{n-2s-4} \text{trace} \left(\left(\sigma^{(1)} \right)^2 - |\xi|^2 \sum_j \left(\sum_i \xi_i \sigma_{ij}^{(1)} \right)^2 \right).$$

The term $u_s^{(3)}$ is computed from (2.7) and (2.8) and represents the interaction of $\sum_{i,j} h_{ij} \partial_{ij}^2$ with the other terms after subtracting the terms which go into $u_s^{(1)}$:

$$(2.19) \quad \begin{aligned} \langle h, (u_s^{(3)}(x, \xi)h) \rangle &= (2S-1)|\xi|^{n-2s-4} \left((\xi \cdot h \cdot \xi) \operatorname{trace} \left(-\sigma^{(1)} - 2\sigma^{(0)} \right) \right. \\ &\quad + |\xi|^2 (\operatorname{tr} h) \operatorname{trace} \left(-\sigma^{(1)} + 2\sigma^{(0)} \right) \\ &\quad \left. + |\xi|^2 \operatorname{trace} \sum_j \sigma_{ij}^{(1)} (\xi_i (h \cdot \xi)_j + \xi_j (h \cdot \xi)_i) \right). \end{aligned}$$

The term $u_s^{(4)}$ is computed from (2.9) and represents the interaction of $\sum_{i,j} h_{ij} \partial_{ij}^2$ with itself after $u_s^{(1)}$ has been taken into account:

$$(2.20) \quad \begin{aligned} \langle h, (u_s^{(4)}(x, \xi)h) \rangle &= \dim E |\xi|^{n-2s-4} \\ &\times \left(\frac{1}{2} |\xi|^4 |h|^2 - 2S |\xi|^2 |h \cdot \xi|^2 + (S+1/4)(\xi \cdot h \cdot \xi)^2 + (S-1)|\xi|^2 (\operatorname{tr} h)(\xi \cdot h \cdot \xi) + \frac{1}{4} |\xi|^4 (\operatorname{tr} h)^2 \right) \\ &= \dim E |\xi|^{n-2s-4} (S-1/2) (-2|\xi|^2 |h \cdot \xi|^2 + (\xi \cdot h \cdot \xi)^2 + |\xi|^2 (\xi \cdot h \cdot \xi)(\operatorname{tr} h)) \\ &\quad + \dim E |\xi|^{n-2s} \left(\frac{1}{2} \operatorname{tr} ((H \Pi_\xi^\perp)^2) + \frac{1}{4} (\operatorname{tr} H \Pi_\xi^\perp)^2 \right). \end{aligned}$$

The fact that (2.11) holds may be verified by explicitly computing $\langle h, u_s(x, \xi)h \rangle$ by substituting (2.14) into (2.3).

3. Calculation of Δ'_p .

To apply these formulas to the Bochner and de Rham Laplacians Δ_p , we must compute the terms of Δ'_p of total degree 2 in the form (2.14). Let $(x^1, \dots, x^n)$ be local coordinates on M , and let $\Lambda^p(M)$ denote the bundle of alternating $(0, p)$ tensors on M . Then the subsets $I \subset \{1, 2, \dots, n\}$ of size p index a basis for $\Lambda_x^p(M)$. The I th basis element is $dx^I = dx^{i_1} \wedge \dots \wedge dx^{i_p}$, where $i_1, \dots, i_p$ are chosen so that $i_1 < i_2 < \dots < i_p$ and $I = \{i_1, \dots, i_p\}$. For a p -form F , we write in components

$$F = \sum_{i_1, \dots, i_p=1}^n F_{i_1 \dots i_p} dx^{i_1} \otimes \dots \otimes dx^{i_p} = \sum_{i_1 < \dots < i_p} F_{i_1 \dots i_p} dx^{i_1} \wedge \dots \wedge dx^{i_p} = \sum_I F_I dx^I.$$

where $F_I := F_{i_1 \dots i_p}$, with $i_1, \dots, i_p$ defined so that $i_1 < \dots < i_p$ and $I = \{i_1, \dots, i_p\}$. It is convenient to write δ for d^* .

Lemma 3.1. *In coordinates which are normal at a point x ,*

$$(3.1) \quad \begin{aligned} ((\nabla^* \nabla)' F)_{i_1 \dots i_p} &\sim \sum_{j,k} h_{jk} \partial_{jk}^2 F_{i_1 \dots i_p} + \sum_{j,k} \left(\partial_j h_{jk} - \frac{1}{2} \partial_k h_{jj} \right) \partial_k F_{i_1 \dots i_p} \\ &+ \sum_{j,k} \sum_{s=1}^p (-1)^{s+1} (\partial_{i_s} h_{jk} + \partial_j h_{i_s k} - \partial_k h_{i_s j}) \partial_j F_{k i_1 \dots \widehat{i_s} \dots i_p} \\ &+ \frac{1}{2} \sum_{j,k} \sum_{s=1}^p (-1)^{s+1} (\partial_{j i_s}^2 h_{jk} - \partial_{jk}^2 h_{i_s j} + \partial_{jj}^2 h_{i_s k}) F_{k i_1 \dots \widehat{i_s} \dots i_p}, \end{aligned}$$

and

$$\begin{aligned}
((\delta' d + d\delta')F)_{i_1 \dots i_p} &\sim \sum_{j,k} h_{jk} \partial_{jk}^2 F_{i_1 \dots i_p} + \sum_{j,k} \left(\partial_j h_{jk} - \frac{1}{2} \partial_k h_{jj} \right) \partial_k F_{i_1 \dots i_p} \\
&+ \sum_{j,k} \sum_{s=1}^p (-1)^{s+1} (\partial_{i_s} h_{jk} + \partial_j h_{i_s k} - \partial_k h_{i_s j}) \partial_j F_{k i_1 \dots \widehat{i_s} \dots i_p} \\
&+ \sum_{j,k} \sum_{s=1}^p (-1)^{s+1} \left(\partial_{i_s j}^2 h_{jk} - \frac{1}{2} \partial_{i_s k}^2 h_{jj} \right) F_{k i_1 \dots \widehat{i_s} \dots i_p} \\
(3.2) \quad &+ \sum_{j,k} \sum_{s=1}^p \sum_{t \neq s} (-1)^{s+t+1} (\partial_{i_s j}^2 h_{i_t k}) F_{j k i_1 \dots \widehat{i_t} \dots \widehat{i_s} \dots i_p}.
\end{aligned}$$

Proof. To carry out this calculation, it is convenient to use the Einstein summation convention that a repeated index in the upper and lower position is summed over. Then

$$(3.3) \quad \nabla^* \nabla F_{i_1 \dots i_p} = -F_{i_1 \dots i_p; j}^j = -g^{j\ell} \partial_\ell F_{i_1 \dots i_s; j} + g^{j\ell} \sum_{s=1}^p (-1)^{s+1} \Gamma_{\ell i_s}^k F_{k i_1 \dots \widehat{i_s} \dots i_p; j} + g^{j\ell} \Gamma_{\ell j}^k F_{i_1 \dots i_s; k}$$

where the Christoffel symbols Γ_{ij}^k are defined by

$$(3.4) \quad \Gamma_{ij}^k = \frac{g^{k\ell}}{2} (\partial_i g_{j\ell} + \partial_j g_{i\ell} - \partial_\ell g_{ij}).$$

Now using the fact that $\Gamma_{ij}^{k'}$ is tensorial in h , and computing in normal coordinates for g , we see that

$$(3.5) \quad G_i^{jk} := g^{j\ell} (\Gamma_{\ell i}^{k'}) = \frac{1}{2} (h_{i; j}^{k j} + h^{jk}_{; i} - h_{i; k}^j),$$

and so

$$\begin{aligned}
&((\nabla^* \nabla)' F)_{i_1 \dots i_p} \\
&\sim h^{j\ell} \partial_\ell F_{i_1 \dots i_s; j} - g^{j\ell} \partial_\ell (F_{i_1 \dots i_s; j}') + \sum_{s=1}^p (-1)^{s+1} G_{i_s}^{jk} \partial_j F_{k i_1 \dots \widehat{i_s} \dots i_p} + G_j^{jk} \partial_k F_{i_1 \dots i_p} \\
&\sim h^{j\ell} \partial_\ell F_{i_1 \dots i_s; j} + g^{j\ell} \partial_\ell \sum_{s=1}^p (-1)^{s+1} \Gamma_{j i_s}^{k'} F_{k i_1 \dots \widehat{i_s} \dots i_p} + \sum_{s=1}^p (-1)^{s+1} G_{i_s}^{jk} \partial_j F_{k i_1 \dots \widehat{i_s} \dots i_p} + G_j^{jk} \partial_k F_{i_1 \dots i_s} \\
&\sim h^{j\ell} \partial_\ell \partial_j F_{i_1 \dots i_s} + 2 \sum_{s=1}^p (-1)^{s+1} G_{i_s}^{jk} \partial_j F_{k i_1 \dots \widehat{i_s} \dots i_p} + G_j^{jk} \partial_k F_{i_1 \dots i_s} + \sum_{s=1}^p (-1)^{s+1} (\partial_j G_{i_s}^{jk}) F_{k i_1 \dots \widehat{i_s} \dots i_p}
\end{aligned}$$

Substituting the formula for G in (3.5) gives (3.1).

For the de Rham Laplacian, we have

$$(3.6) \quad (dF)_{i_1 \dots i_{p+1}} = \sum_{s=1}^{p+1} (-1)^{s+1} \partial_{i_s} F_{i_1 \dots \widehat{i_s} \dots i_{p+1}},$$

and

$$(3.7) \quad (\delta F)_{i_1 \dots i_{p-1}} = -F_{j i_1 \dots i_{p-1}}{}^j.$$

But

$$(3.8) \quad -F_{j i_1 \dots i_{p-1}}{}^j = -g^{j\ell} F_{j i_1 \dots i_{p-1}; \ell} \\ = g^{j\ell} \left(-\partial_\ell F_{j i_1 \dots i_{p-1}} + \Gamma_{\ell j}^k F_{k i_1 \dots i_{p-1}} + \sum_{s=1}^{p-1} (-1)^{s+1} \Gamma_{\ell i_s}^k F_{j k i_1 \dots \widehat{i_s} \dots i_{p-1}} \right),$$

and so from (3.5),

$$(3.9) \quad \left(-F_{j i_1 \dots i_{p-1}}{}^j \right)' = h^{jk} F_{j i_1 \dots i_{p-1}; k} + G_j^{jk} F_{k i_1 \dots i_{p-1}} + \sum_{s=1}^{p-1} (-1)^{s+1} G_{i_s}^{jk} F_{j k i_1 \dots \widehat{i_s} \dots i_{p-1}}.$$

Set

$$(3.10) \quad 1_{t>s} = \begin{cases} 1 & t > s \\ 0 & t \leq s \end{cases}.$$

Then

$$(3.11) \quad (\delta' dF)_{i_1 \dots i_p} = \left(-(dF)_{j i_1 \dots i_p}{}^j \right)' \\ \sim h^{jk} \partial_k (dF)_{j i_1 \dots i_p} + G_j^{jk} (dF)_{k i_1 \dots i_p} + \sum_{s=1}^p (-1)^{s+1} G_{i_s}^{jk} (dF)_{j k i_1 \dots \widehat{i_s} \dots i_p} \\ = h^{jk} \partial_{jk}^2 F_{i_1 \dots i_p} + \sum_{s=1}^p (-1)^s h^{jk} \partial_{k i_s}^2 F_{j i_1 \dots \widehat{i_s} \dots i_p} \\ + G_j^{jk} \partial_k F_{i_1 \dots i_p} + G_j^{jk} \sum_{s=1}^p (-1)^s \partial_{i_s} F_{k i_1 \dots \widehat{i_s} \dots i_p} \\ + \sum_{s=1}^p (-1)^{s+1} \left(G_{i_s}^{jk} - G_{i_s}^{kj} \right) \partial_j F_{k i_1 \dots \widehat{i_s} \dots i_p} \\ + \sum_{s=1}^p \sum_{t \neq s} (-1)^{s+t+1_{t>s}} G_{i_s}^{jk} \partial_{i_t} F_{j k i_1 \dots \widehat{i_t} \dots \widehat{i_s} \dots i_p}.$$

On the other hand,

$$(d\delta' F)_{i_1 \dots i_p} = \sum_{s=1}^p (-1)^{s+1} \partial_{i_s} (\delta' F)_{i_1 \dots \widehat{i_s} \dots i_p} \\ = \sum_{s=1}^p (-1)^{s+1} \partial_{i_s} \left(h^{jk} F_{j i_1 \dots \widehat{i_s} \dots i_p; k} + G_j^{jk} F_{k i_1 \dots \widehat{i_s} \dots i_p} + \sum_{t \neq s} (-1)^{t+1_{s>t}} G_{i_t}^{jk} F_{j k i_1 \dots \widehat{i_t} \dots \widehat{i_s} \dots i_p} \right)$$

$$\begin{aligned}
& \sim \sum_{s=1}^p (-1)^{s+1} (\partial_{i_s} h^{jk}) \partial_k F_{j i_1 \dots \widehat{i_s} \dots i_p} + \sum_{s=1}^p (-1)^{s+1} h^{jk} \partial_{i_s k}^2 F_{j i_1 \dots \widehat{i_s} \dots i_p} \\
& + \sum_{s=1}^p (-1)^{s+1} (\partial_{i_s} G_j^{jk}) F_{k i_1 \dots \widehat{i_s} \dots i_p} + \sum_{s=1}^p (-1)^{s+1} G_j^{jk} \partial_{i_s} F_{k i_1 \dots \widehat{i_s} \dots i_p} \\
(3.12) \quad & + \sum_{s=1}^p \sum_{t \neq s} (-1)^{s+t+1} {}_{t>s} \left(\partial_{i_s} G_{i_t}^{jk} \right) F_{j k i_1 \dots \widehat{i_t} \dots \widehat{i_s} \dots i_p} + \sum_{s=1}^p \sum_{t \neq s} (-1)^{s+t+1} {}_{t>s} G_{i_t}^{jk} \partial_{i_s} F_{j k i_1 \dots \widehat{i_t} \dots \widehat{i_s} \dots i_p}
\end{aligned}$$

We notice that the second terms on the last three lines of (3.12) cancel with terms in (3.11), and hence

$$\begin{aligned}
((\delta' d + d \delta') F)_{i_1 \dots i_p} & \sim h^{jk} \partial_{j k}^2 F_{i_1 \dots i_p} + G_j^{jk} \partial_k F_{i_1 \dots i_p} \\
& + \sum_{s=1}^p (-1)^{s+1} (\partial_{i_s} h^{jk}) \partial_k F_{j i_1 \dots \widehat{i_s} \dots i_p} + \sum_{s=1}^p (-1)^{s+1} (G_{i_s}^{jk} - G_{i_s}^{kj}) \partial_j F_{k i_1 \dots \widehat{i_s} \dots i_p} \\
& + \sum_{s=1}^p (-1)^{s+1} (\partial_{i_s} G_j^{jk}) F_{k i_1 \dots \widehat{i_s} \dots i_p} + \sum_{s=1}^p \sum_{t \neq s} (-1)^{s+t+1} {}_{t>s} (\partial_{i_s} G_{i_t}^{jk}) F_{j k i_1 \dots \widehat{i_t} \dots \widehat{i_s} \dots i_p}
\end{aligned}$$

which gives (3.2) on substituting in the formula for G in (3.5).

4. Matrix entries of Δ'_p .

As we already discussed at the beginning of Section 3, if we fix local coordinates on M about the point x , then dx^I forms a basis for Λ_x^p , where I ranges over subsets of $\{1, 2, \dots, n\}$ of size p . A p -form is then identified with a vector valued function (F_I) , and the Bochner or de Rham Laplacian at x can be identified with a matrix of (scalar) differential operators. The purpose of this section is to identify these matrix entries, or rather the terms of total degree 2 in these entries (see (2.13)). We will start by considering the Bochner Laplacian $\nabla^* \nabla$, whose terms of total degree 2 are given in (3.1), which we rewrite here:

$$\begin{aligned}
((\nabla^* \nabla)' F)_{i_1 \dots i_p} & \sim \sum_{j,k} h_{jk} \partial_{j k}^2 F_{i_1 \dots i_p} + \sum_{j,k} \left(\partial_j h_{jk} - \frac{1}{2} \partial_k h_{jj} \right) \partial_k F_{i_1 \dots i_p} \\
& + \sum_{j,k} \sum_{s=1}^p (-1)^{s+1} (\partial_{i_s} h_{jk} + \partial_j h_{i_s k} - \partial_k h_{i_s j}) \partial_j F_{k i_1 \dots \widehat{i_s} \dots i_p} \\
(4.1) \quad & + \frac{1}{2} \sum_{j,k} \sum_{s=1}^p (-1)^{s+1} (\partial_{j i_s}^2 h_{jk} - \partial_{j k}^2 h_{i_s j} + \partial_{j j}^2 h_{i_s k}) F_{k i_1 \dots \widehat{i_s} \dots i_p},
\end{aligned}$$

Suppose that $I = \{i_1, \dots, i_p\}$ where $i_1 < \dots < i_p$. The terms on the top line of (4.1) give II matrix entries. The summands appearing in the second and third lines give KI matrix entries, where $K = (I \setminus \{i_s\}) \cup \{k\}$. We get $K = I$ when $k = i_s$, so altogether, the II matrix entry of (4.1) is

$$(4.2) \quad \sum_{j,k} h_{jk} \partial_{j k}^2 + \sum_{j,k} \left(\partial_j h_{jk} - \frac{1}{2} \partial_k h_{jj} \right) \partial_k + \sum_j \sum_{i \in I} (\partial_j h_{ii}) \partial_j + \frac{1}{2} \sum_j \sum_{i \in I} (\partial_{jj}^2 h_{ii}).$$

It is convenient to introduce the notation

$$(4.3) \quad \chi_I(j) = \begin{cases} 1 & j \in I, \\ 0 & j \notin I, \end{cases}$$

$$(4.4) \quad \text{sgn}_I(j) = \begin{cases} 1 & j \in I, \\ -1 & j \notin I. \end{cases}$$

After simplifying (4.2), we get

Lemma 4.1. *The term of total degree 2 in the II matrix entry of $\nabla^* \nabla$ is*

$$(4.5) \quad \sum_{i,j} (h_{ij} \partial_{ij}^2 + \partial_i h_{ij} \partial_j) + \frac{1}{2} \sum_{i,j} \text{sgn}_I(i) \partial_j h_{ii} \partial_j + \frac{1}{2} \sum_{i,j} \chi_I(i) (\partial_{jj}^2 h_{ii}).$$

The only other non-vanishing matrix entries of total degree 2 in $\nabla^* \nabla$ are KI terms where $K = (I \setminus \{i_s\}) \cup \{k\}$, and $k \notin I$. These can easily be read off from (4.1), although one has to pay attention to the sign.

Definition 4.2. For $J \subset \{1, 2, \dots, n\}$ and $i \in \{1, 2, \dots, n\}$, define

$$(4.6) \quad N(i, J) = \#\{j \in J : j < i\}.$$

The point of this definition is that

$$(4.7) \quad dx^{\{i\} \cup J} = (-1)^{N(i, J)} dx^i \wedge dx^J.$$

Lemma 4.3. *Suppose that J is a subset of $\{1, \dots, n\}$ of size $p-1$, and $I = J \cup \{i\}$ and $K = J \cup \{k\}$, where $i, k \notin J$ and $i \neq k$. Then the term of total degree 2 in the KI matrix entry of $\nabla^* \nabla$ is*

$$(4.8) \quad (-1)^{N(i, J) + N(k, J)} \sum_j \left((\partial_i h_{jk} + \partial_j h_{ik} - \partial_k h_{ij}) \partial_j + \frac{1}{2} (\partial_{ij}^2 h_{jk} - \partial_{jk}^2 h_{ij} + \partial_{jj}^2 h_{ik}) \right).$$

We can carry out the same analysis for the de Rham Laplacian in (3.2). The only new feature is a new type of term appearing in the last line of (3.2), which we rewrite here:

$$(4.9) \quad \begin{aligned} ((\delta' d + d\delta')F)_{i_1 \dots i_p} &\sim \dots\dots\dots \\ &+ \sum_{s=1}^p \sum_{t \neq s} \sum_{k, \ell} (-1)^{s+t+1_{t>s}} (\partial_{isk}^2 h_{it\ell}) F_{k\ell i_1 \dots \widehat{i_t} \dots \widehat{i_s} \dots i_p}. \end{aligned}$$

By writing $J = I \setminus \{i_s, i_t\}$ and $L = J \cup \{k, \ell\}$ and noting that

$$F_{i_1 \dots i_p} = (-1)^{s+t+1_{t>s}} F_{i_s i_t i_1 \dots \widehat{i_s} \dots \widehat{i_t} \dots i_p},$$

we see that LI matrix entry in the summand of (4.9) is

$$(4.10) \quad (-1)^{N(i_s, J) + N(i_t, J) + N(k, J) + N(\ell, J)} \operatorname{sgn}((i_t - i_s)(\ell - k)) (\partial_{i_s k}^2 h_{i_t \ell}).$$

The case $L = I$ can occur in two ways: $k = i_s$ and $k = i_t$, or $j = i_t$ and $k = i_s$. Hence after carrying out the k, ℓ sum (but not the s, t sum), the II matrix entry of (4.9) is

$$(4.11) \quad (\partial_{i_s i_s}^2 h_{i_t i_t}) - (\partial_{i_s i_t}^2 h_{i_t i_s}).$$

Hence the II matrix entry of (3.2) is

$$(4.12) \quad \sum_{j, k} h_{jk} \partial_{jk}^2 + \sum_{j, k} \left(\partial_j h_{jk} - \frac{1}{2} \partial_k h_{jj} \right) \partial_k \\ + \sum_{i, j} \chi_I(i) \left((\partial_j h_{ii}) \partial_j + \partial_{ij}^2 h_{ij} - \frac{1}{2} \partial_{ii}^2 h_{jj} \right) + \sum_{i, j} \chi_I(i) \chi_{I'}(j) (\partial_{ii}^2 h_{jj} - \partial_{ij}^2 h_{ij}).$$

Simplifying this, we get the following.

Lemma 4.4. *The term of total degree 2 in the II matrix entry of $d^*d + dd^*$ is*

$$(4.13) \quad \sum_{i, j} (h_{ij} \partial_{ij}^2 + \partial_i h_{ij} \partial_j) + \frac{1}{2} \sum_{i, j} \operatorname{sgn}_I(j) \partial_i h_{jj} \partial_i \\ + \sum_{i, j} \chi_I(i) \chi_{I'}(j) (\partial_{ij}^2 h_{ij}) + \frac{1}{2} \sum_{i, j} \chi_I(i) \operatorname{sgn}_I(j) \partial_{ii}^2 h_{jj}.$$

Now let $\bar{J}$ be a subset of $\{1, \dots, n\}$ of size $p-1$ and let p, q with $p \neq q$ be such that $p \notin \bar{J}$ and $q \notin \bar{J}$. Set $I = \bar{J} \cup \{p\}$ and $K = \bar{J} \cup \{q\}$. The the summand in (4.9) gives an LI matrix entry, and $L = K$ if one of the following four cases holds:

$$(4.14) \quad \begin{array}{lll} i_s = p, & k = q, & i_t = \ell, \\ i_t = p, & k = q, & i_s = \ell, \end{array} \quad \begin{array}{lll} i_s = p, & \ell = q, & i_t = k, \\ i_t = p, & \ell = q, & i_s = k. \end{array}$$

Hence the KI matrix entry of (4.9) after summing is

$$(4.15) \quad (-1)^{N(p, \bar{J}) + N(q, \bar{J})} \sum_{j \in I} (\partial_{pq}^2 h_{jj} + \partial_{jj}^2 h_{pq} - \partial_{jp}^2 h_{jq} - \partial_{jq}^2 h_{jp}).$$

In order to see this, it is important to note that the sets J from (4.10) and $\bar{J}$ differ by one element, i_s or i_t . By considering the other terms in (3.2), we get the following.

Lemma 4.5. *For $I = J \cup \{i\}$ and $K = J \cup \{k\}$, where $i, k \notin J$ and $i \neq k$, the term of total degree 2 in the KI matrix entry of $d^*d + dd^*$ is*

$$(4.16) \quad (-1)^{N(i, J) + N(k, J)} \left(\sum_j ((\partial_i h_{jk} + \partial_j h_{ik} - \partial_k h_{ij}) \partial_j + (\partial_{ij}^2 h_{jk}) - \frac{1}{2} (\partial_{ik}^2 h_{jj})) \right. \\ \left. + \sum_j \chi_I(j) (\partial_{ik}^2 h_{jj} + \partial_{jj}^2 h_{ik} - \partial_{ij}^2 h_{jk} - \partial_{jk}^2 h_{ij}) \right)$$

Finally, similar considerations as in the previous case give the following.

Lemma 4.6. Suppose J is a subset of $\{1, \dots, n\}$ of size $p-2$, and $I = J \cup \{i, j\}$, $L = J \cup \{k, \ell\}$, where i, j, k, ℓ are distinct with $i < j$, $k < \ell$ and $i, j, k, \ell \notin J$. Then the term of total degree 2 in the LI matrix entry of $d^*d + dd^*$ is

$$(4.16) \quad (-1)^{N(i,J)+N(j,J)+N(k,J)+N(\ell,J)} \left(\partial_{ik}^2 h_{j\ell} + \partial_{j\ell}^2 h_{ik} - \partial_{i\ell}^2 h_{jk} - \partial_{jk}^2 h_{i\ell} \right).$$

Because i, j, k, ℓ are all distinct, we see that this vanishes if h is diagonal.

5. Proof of Theorem 1 for the Bochner Laplacian.

Combining the following Lemma with (2.12) and (2.20), gives Theorem 1 for the Bochner Laplacian, see (1.7), (1.8) and (1.9). (Here in the proof we are dealing with the real case. The factor 2 in (1.7) is the dimension of the complex line.)

Lemma 5.1. For the Bochner Laplacian $\nabla^* \nabla$ on p forms,

$$(5.1) \quad \langle h, u_s^{(1)}(x, \xi) h \rangle = (S^2 - 1/4) |\xi|^{n-2s} \binom{n}{p} (\text{tr}(H \Pi_\xi^\perp))^2.$$

$$(5.2) \quad \begin{aligned} \langle h, u^{(2)}(x, \xi) h \rangle &= (S - 1/2) |\xi|^{n-2s} 4 \binom{n-2}{p-1} \text{tr}((H \Pi_\xi^\perp)^2) \\ &\quad + (S - 1/2) |\xi|^{n-2s-4} \binom{n}{p} (-2|\xi|^2 |h \cdot \xi|^2 + 2(\xi \cdot h \cdot \xi)^2). \end{aligned}$$

$$(5.3) \quad \begin{aligned} \langle h, u^{(3)}(x, \xi) h \rangle &= (S - 1/2) |\xi|^{n-2s} \binom{n}{p} (\text{tr}(H \Pi_\xi^\perp))^2 \\ &\quad + (S - 1/2) |\xi|^{n-2s-4} \binom{n}{p} (4|\xi|^2 |h \cdot \xi|^2 - 3(\xi \cdot h \cdot \xi)^2 - |\xi|^2 (\xi \cdot h \cdot \xi) (\text{tr } h)). \end{aligned}$$

Proof. For $j = 0, 1, 2$, and subsets I, K of $\{1, \dots, n\}$ of size p , write $\sigma^{(j)KI}$ for the KI matrix element of $\sigma^{(j)}$, and write $\sigma_{pq}^{(j)KI}$ for the KI matrix element of $\sigma_{pq}^{(j)}$. Now for a general operator on p -forms, we can expand the trace in (2.17) to get

$$(5.4) \quad \begin{aligned} \langle h, u_s^{(1)} h \rangle &= (S^2 - 1/4) |\xi|^{n-2s-4} \left(\sum_I \left(\sigma^{(2)II} - 2\sigma^{(1)II} + 4\sigma^{(0)II} \right)^2 \right. \\ &\quad \left. + \sum_{K \neq I} \left(\sigma^{(2)KI} - 2\sigma^{(1)KI} + 4\sigma^{(0)KI} \right) \left(\sigma^{(2)IK} - 2\sigma^{(1)IK} + 4\sigma^{(0)IK} \right) \right). \end{aligned}$$

For the Bochner Laplacian, we see from Lemma 4.1, we see that

$$(5.5) \quad \begin{aligned} \sigma^{(2)II} &= \xi \cdot h \cdot \xi, & \sigma^{(1)II} &= (\xi \cdot h \cdot \xi) + \frac{1}{2} |\xi|^2 \sum_i \text{sgn}_I(i) h_{ii}, & \sigma^{(0)II} &= \frac{1}{2} |\xi|^2 \sum_i \chi_I(i) h_{ii}. \end{aligned}$$

Hence

$$(5.6) \quad \sigma^{(2)II} - 2\sigma^{(1)II} + 4\sigma^{(0)II} = -(\xi \cdot h \cdot \xi) + |\xi|^2(\text{tr } h).$$

On the other hand, let I, J, K be as in Lemma 4.3. Then

$$(5.7) \quad \begin{aligned} \sigma^{(1)KI} &= (-1)^{N(i,J)+N(k,J)} (\xi_i(h \cdot \xi)_k - \xi_k(h \cdot \xi)_i + |\xi|^2 h_{ik}), \\ \sigma^{(0)KI} &= (-1)^{N(i,J)+N(k,J)} \frac{1}{2} (\xi_i(h \cdot \xi)_k - \xi_k(h \cdot \xi)_i + |\xi|^2 h_{ik}). \end{aligned}$$

We see that

$$(5.8) \quad \sigma^{(2)KI} - 2\sigma^{(1)KI} + 4\sigma^{(0)KI} = 0.$$

Hence from (5.6) and (5.8), we see that for the Bochner Laplacian,

$$(5.9) \quad \begin{aligned} \langle h, u_s^{(1)} h \rangle &= (S^2 - 1/4) |\xi|^{n-2s-4} \sum_I (-(\xi \cdot h \cdot \xi) + |\xi|^2(\text{tr } h))^2 \\ &= (S^2 - 1/4) |\xi|^{n-2s-4} \binom{n}{p} (-(\xi \cdot h \cdot \xi) + |\xi|^2(\text{tr } h))^2, \end{aligned}$$

which gives (5.1).

Now we prove (5.2). From (2.18), we can write

$$(5.10) \quad \begin{aligned} \langle h, u_s^{(2)} h \rangle &= (2S - 1) |\xi|^{n-2s-4} \left(\sum_I \left((\sigma^{(1)II})^2 - |\xi|^2 \sum_j \left(\sum_i \xi_i \sigma_{ij}^{(1)II} \right)^2 \right) \right. \\ &\quad \left. + \sum_{I \neq K} \left(\sigma^{(1)KI} \sigma^{(1)IK} - |\xi|^2 \sum_q \left(\sum_p \xi_p \sigma_{pq}^{(1)KI} \right) \left(\sum_r \xi_r \sigma_{rq}^{(1)IK} \right) \right) \right). \end{aligned}$$

Now from (5.5) and (5.7) we have

$$(5.11) \quad (\sigma^{(1)II})^2 = (\xi \cdot h \cdot \xi)^2 + |\xi|^2 (\xi \cdot h \cdot \xi) \sum_i \text{sgn}_I(i) h_{ii} + \frac{1}{4} |\xi|^4 \left(\sum_i \text{sgn}_I(i) h_{ii} \right)^2.$$

and with I, K as in Lemma 4.3,

$$(5.12) \quad \begin{aligned} \sigma^{(1)KI} \sigma^{(1)IK} &= (\xi_i(h \cdot \xi)_k - \xi_k(h \cdot \xi)_i + |\xi|^2 h_{ik}) (\xi_k(h \cdot \xi)_i - \xi_i(h \cdot \xi)_k + |\xi|^2 h_{ik}) \\ &= -(\xi_i(h \cdot \xi)_k - \xi_k(h \cdot \xi)_i)^2 + |\xi|^4 h_{ik}^2. \end{aligned}$$

With I, J, K as in Lemma 4.3, we see from Lemmas 4.1 and 4.3, that

$$(5.13) \quad \begin{aligned} \sigma_{pq}^{(1)II} &= h_{pq} + \frac{\delta_{pq}}{2} \sum_i \text{sgn}_I(i) h_{ii}, & \sum \xi_p \sigma_{pq}^{(1)II} &= (h \cdot \xi)_q + \frac{1}{2} \xi_q \left(\sum_i \text{sgn}_I(i) h_{ii} \right), \\ \sigma_{pq}^{(1)KI} &= \delta_{ip} h_{qk} - \delta_{kp} h_{iq} + \delta_{pq} h_{ik}, & \sum \xi_p \sigma_{pq}^{(1)KI} &= \xi_i h_{qk} - \xi_k h_{iq} + \xi_q h_{ik}. \end{aligned}$$

We see that

$$(5.14) \quad |\xi|^2 \sum_j \left(\sum_i \xi_i \sigma_{ij}^{(1)II} \right)^2 = |\xi|^2 |h \cdot \xi|^2 + |\xi|^2 (\xi \cdot h \cdot \xi) \sum_i \operatorname{sgn}_I(i) h_{ii} + \frac{1}{4} |\xi|^4 \left(\sum_i \operatorname{sgn}_I(i) h_{ii} \right)^2,$$

and

$$(5.15) \quad |\xi|^2 \sum_q \left(\sum_p \xi_p \sigma_{pq}^{(1)KI} \right) \left(\sum_r \xi_r \sigma_{rq}^{(1)IK} \right) = -|\xi|^2 \sum_q (\xi_i h_{kq} - \xi_k h_{iq})^2 + |\xi|^4 h_{ik}^2.$$

Putting (5.11), (5.12), (5.14) and (5.15) into (5.10), we get

$$(5.16) \quad \langle h, u^{(2)} h \rangle = |\xi|^{n-2s-4} (2S-1) \left(\sum_I ((\xi \cdot h \cdot \xi)^2 - |\xi|^2 |h \cdot \xi|^2) \right. \\ \left. - \sum_I \sum_{i,k} \chi_I(i) \chi_{I'}(k) \left((\xi_i (h \cdot \xi)_k - \xi_k (h \cdot \xi)_i)^2 - |\xi|^2 \sum_q (\xi_i h_{kq} - \xi_k h_{iq})^2 \right) \right).$$

Here, we use the fact that we can sum over all I and K of the form in Lemma 4.3, by summing over I , and over those i, k with $i \in I$ and $k \notin I$. Using (7.1), we find that (5.16) equals

$$(5.17) \quad |\xi|^{n-2s-4} (2S-1) \left(\binom{n}{p} ((\xi \cdot h \cdot \xi)^2 - |\xi|^2 |h \cdot \xi|^2) \right. \\ \left. + \binom{n-2}{p-1} \sum_{i,k} \left(|\xi|^2 \sum_q (\xi_i h_{kq} - \xi_k h_{iq})^2 - (\xi_i (h \cdot \xi)_k - \xi_k (h \cdot \xi)_i)^2 \right) \right).$$

This equals (5.2).

Now from (2.19),

$$(5.18) \quad \langle h, u_s^{(3)} h \rangle = (2S-1) |\xi|^{n-2s-4} \sum_I \left((\xi \cdot h \cdot \xi) \left(-\sigma^{(1)II} - 2\sigma^{(0)II} \right) \right. \\ \left. + |\xi|^2 (\operatorname{tr} h) \left(-\sigma^{(1)II} + 2\sigma^{(0)II} \right) + |\xi|^2 \sum_j \sigma_{ij}^{(1)II} (\xi_i (h \cdot \xi)_j + \xi_j (h \cdot \xi)_i) \right)$$

Using (5.5) and (5.13), for the Bochner Laplacian we get

$$-\sigma^{(1)II} - 2\sigma^{(0)II} = -(\xi \cdot h \cdot \xi) - |\xi|^2 \sum_i \left(\frac{1}{2} + \operatorname{sgn}_I(i) \right) h_{ii} \\ -\sigma^{(1)II} + 2\sigma^{(0)II} = -(\xi \cdot h \cdot \xi) + \frac{1}{2} |\xi|^2 (\operatorname{tr} h) \\ \sum_j \sigma_{ij}^{(1)II} (\xi_i (h \cdot \xi)_j + \xi_j (h \cdot \xi)_i) = 2|h \cdot \xi|^2 + (\xi \cdot h \cdot \xi) \sum_i \operatorname{sgn}_I(i) h_{ii}.$$

Hence putting this into (5.18),

$$\begin{aligned} \langle h, u_s^{(3)} h \rangle &= (2S-1)|\xi|^{n-2s-4} \sum_I \left((\xi \cdot h \cdot \xi) \left(-(\xi \cdot h \cdot \xi) - |\xi|^2 \sum_i \left(\frac{1}{2} + \text{sgn}_I(i) \right) h_{ii} \right) \right. \\ &\quad \left. + |\xi|^2 (\text{tr } h) \left(-(\xi \cdot h \cdot \xi) + \frac{1}{2} |\xi|^2 (\text{tr } h) \right) + |\xi|^2 \left(2|h \cdot \xi|^2 + (\xi \cdot h \cdot \xi) \sum_i \text{sgn}_I(i) h_{ii} \right) \right) \end{aligned}$$

The terms involving $\sum_i \text{sgn}_I(i) h_{ii}$ cancel, and we are left with

$$(2S-1)|\xi|^{n-2s-4} \binom{n}{p} \left(2|\xi|^2 |h \cdot \xi|^2 - (\xi \cdot h \cdot \xi)^2 - \frac{3}{2} |\xi|^2 (\xi \cdot h \cdot \xi) (\text{tr } h) + \frac{1}{2} |\xi|^4 (\text{tr } h)^2 \right),$$

which equals (5.3).

6. Proof of Theorem 1 for the de Rham Laplacian.

Combining the following Lemma with (2.12) and (2.20), gives Theorem 1 for the de Rham Laplacian, see (1.7), (1.10) and (1.11). (Here in the proof we are dealing with the real case. The factor 2 in (1.7) is the dimension of the complex line.)

Lemma 5.2. *For the de Rham Laplacian on p forms,*

$$\begin{aligned} \langle h, u_s^{(1)} h \rangle &= (S^2 - 1/4) |\xi|^{n-2s} \left(4 \binom{n-2}{p-1} \text{tr}((H\Pi_\xi^\perp)^2) + \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) (\text{tr}(H\Pi_\xi^\perp))^2 \right). \\ \langle h, u_s^{(2)} h \rangle &= (S-1/2) |\xi|^{n-2s} 4 \binom{n-2}{p-1} \text{tr}((H\Pi_\xi^\perp)^2) \\ &\quad + (S-1/2) |\xi|^{n-2s-4} \binom{n}{p} (-2|\xi|^2 |h \cdot \xi|^2 + 2(\xi \cdot h \cdot \xi)^2). \\ \langle h, u_s^{(3)} h \rangle &= (S-1/2) |\xi|^{n-2s} \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) (\text{tr}(H\Pi_\xi^\perp))^2 \\ &\quad + (S-1/2) |\xi|^{n-2s-4} \binom{n}{p} (4|\xi|^2 |h \cdot \xi|^2 - 3(\xi \cdot h \cdot \xi)^2 - |\xi|^2 (\xi \cdot h \cdot \xi) (\text{tr } h)). \end{aligned}$$

Proof. We work in coordinates ξ in which h is diagonal. This is essential for the calculation of $u_s^{(1)}$, which becomes extremely lengthy if we drop this assumption. We know that $u_s^{(1)}$ is itself independent of the coordinates, because it is the coefficient of u_s in the variable s^2 . To compute $u_s^{(1)}$ we apply (5.4) and use the notation given there. By Lemma 4.4, for the de Rham Laplacian,

$$\begin{aligned} \sigma^{(2)II} &= \xi \cdot h \cdot \xi, \\ \sigma^{(1)II} &= (\xi \cdot h \cdot \xi) + \frac{1}{2} |\xi|^2 \sum_i \text{sgn}_I(i) h_{ii}, \\ (6.4) \quad \sigma^{(0)II} &= \sum_{i,j} \chi_I(i) \chi_{I'}(j) \xi_i \xi_j h_{ij} + \frac{1}{2} \sum_{i,j} \chi_I(j) \text{sgn}_I(i) \xi_j^2 h_{ii} = \frac{1}{2} \sum_{i,j} \chi_I(j) \text{sgn}_I(i) \xi_j^2 h_{ii}, \end{aligned}$$

where the last equality holds since h is diagonal. Hence

$$(6.5) \quad \sigma^{(2)II} - 2\sigma^{(1)II} + 4\sigma^{(0)II} = -(\xi \cdot h \cdot \xi) + \sum_{i,j} \text{sgn}_I(i) \text{sgn}_I(j) \xi_i^2 h_{jj}.$$

On the other hand, let I, J, K be as in Lemma 4.5. Then $\sigma^{(2)KI} = 0$, and

$$(5.7) \quad \begin{aligned} \sigma^{(1)KI} &= (-1)^{N(i,J)+N(k,J)} (\xi_i(h \cdot \xi)_k - \xi_k(h \cdot \xi)_i + |\xi|^2 h_{ik}), \\ \sigma^{(0)KI} &= (-1)^{N(i,J)+N(k,J)} \left(\xi_i(h \cdot \xi)_k - \frac{1}{2} \xi_i \xi_k (\text{tr } h) \right. \\ (6.6) \quad &\quad \left. + \sum_j \chi_I(j) (\xi_i \xi_k h_{jj} + \xi_j^2 h_{ik} - \xi_i \xi_j h_{jk} - \xi_i \xi_k h_{ij}) \right). \end{aligned}$$

We see that

$$(6.7) \quad \begin{aligned} \sigma^{(2)KI} - 2\sigma^{(1)KI} + 4\sigma^{(0)KI} &= (-1)^{N(i,J)+N(k,J)} 2 \sum_j \text{sgn}_I(j) (\xi_i \xi_k h_{jj} + \xi_j^2 h_{ik} - \xi_i \xi_j h_{jk} - \xi_i \xi_k h_{ij}) \\ &= (-1)^{N(i,J)+N(k,J)} 2 \left(\left(\sum_j \text{sgn}_I(j) h_{jj} \right) + h_{kk} - h_{ii} \right) \xi_i \xi_k, \end{aligned}$$

where the last line follows because h is diagonal and $i \in I, k \notin I$. Because of Lemma 4.6, we have now considered all the non-zero matrix entries of the coefficient symbols. From (5.4) we get

$$\begin{aligned} \langle h, u_s^{(1)} h \rangle &= (S^2 - 1/4) |\xi|^{n-2s-4} \sum_I \left(\left(-(\xi \cdot h \cdot \xi) + \sum_{i,j} \text{sgn}_I(i) \text{sgn}_I(j) \xi_i^2 h_{jj} \right)^2 \right. \\ &\quad \left. + 4 \sum_{i,k} \chi_I(i) \chi_{I'}(k) \xi_i^2 \xi_k^2 \left(\left(\sum_j \text{sgn}_I(j) h_{jj} \right)^2 - (h_{kk} - h_{ii})^2 \right) \right) \end{aligned}$$

Multiplying out the brackets, we get

$$(6.8) \quad \begin{aligned} \langle h, u_s^{(1)} h \rangle &= (S^2 - 1/4) |\xi|^{n-2s-4} \left(\binom{n}{p} (\xi \cdot h \cdot \xi)^2 - 2(\xi \cdot h \cdot \xi) \sum_{i,j} \text{sgn}_I(i) \text{sgn}_I(j) \xi_i^2 h_{jj} \right. \\ &\quad \left. + \sum_{i,j,k,\ell} \sum_I \text{sgn}_I(j) \text{sgn}_I(\ell) (\text{sgn}_I(i) \text{sgn}_I(k) + 4\chi_I(i) \chi_{I'}(k)) \xi_i^2 \xi_k^2 h_{jj} h_{\ell\ell} \right. \\ &\quad \left. + 4 \sum_{i,k} \sum_I \chi_I(i) \chi_{I'}(k) \xi_i^2 \xi_k^2 (2h_{ii} h_{kk} - h_{ii}^2 - h_{kk}^2) \right). \end{aligned}$$

But we can considerably simplify the second line of (6.8) because

$$\begin{aligned} \sum_{i,k} (\operatorname{sgn}_I(i) \operatorname{sgn}_I(k) + 4\chi_I(i)\chi_{I'}(k)) \xi_i^2 \xi_k^2 \\ = \sum_{i,k} (\operatorname{sgn}_I(i) \operatorname{sgn}_I(k) + 2\chi_I(i)\chi_{I'}(k) + 2\chi_{I'}(i)\chi_I(k)) \xi_i^2 \xi_k^2 = \sum_{i,k} \xi_i^2 \xi_k^2 = |\xi|^4. \end{aligned}$$

Hence we get

$$\begin{aligned} (6.9) \quad \langle h, u_s^{(1)} h \rangle &= C(s)(S^2 - 1/4)|\xi|^{n-2s-4} \left(\binom{n}{p} (\xi \cdot h \cdot \xi)^2 - 2(\xi \cdot h \cdot \xi) \sum_{i,j} \sum_I \operatorname{sgn}_I(i) \operatorname{sgn}_I(j) \xi_i^2 h_{jj} \right. \\ &\quad \left. + |\xi|^4 \sum_{j,\ell} \sum_I \operatorname{sgn}_I(j) \operatorname{sgn}_I(\ell) h_{jj} h_{\ell\ell} + 4 \sum_{i,k} \sum_I \chi_I(i) \chi_{I'}(k) \xi_i^2 \xi_k^2 (2h_{ii} h_{kk} - h_{ii}^2 - h_{kk}^2) \right). \end{aligned}$$

From (7.1) and (7.2),

$$\begin{aligned} \sum_{i,j} \sum_I \operatorname{sgn}_I(i) \operatorname{sgn}_I(j) \xi_i^2 h_{jj} &= \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) |\xi|^2 (\operatorname{tr} h) + 4 \binom{n-2}{p-1} (\xi \cdot h \cdot \xi), \\ \sum_I \operatorname{sgn}_I(j) \operatorname{sgn}_I(\ell) h_{jj} h_{\ell\ell} &= \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) (\operatorname{tr} h)^2 + 4 \binom{n-2}{p-1} |h|^2, \\ \sum_{i,k} \sum_I \chi_I(i) \chi_{I'}(k) \xi_i^2 \xi_k^2 (2h_{ii} h_{kk} - h_{ii}^2 - h_{kk}^2) &= 2 \binom{n-2}{p-1} ((\xi \cdot h \cdot \xi)^2 - |\xi|^2 |h \cdot \xi|^2). \end{aligned}$$

Hence putting this into (6.9), we get

$$\begin{aligned} \langle h, u_s^{(1)} h \rangle &= (S^2 - 1/4)|\xi|^{n-2s-4} \left(\binom{n}{p} (\xi \cdot h \cdot \xi)^2 - 2 \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) |\xi|^2 (\xi \cdot h \cdot \xi) (\operatorname{tr} h) \right. \\ &\quad \left. + \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) |\xi|^4 (\operatorname{tr} h)^2 + 4 \binom{n-2}{p-1} |\xi|^4 |h|^2 - 8 \binom{n-2}{p-1} |\xi|^2 |h \cdot \xi|^2 \right), \end{aligned}$$

which equals (6.1).

The proof of (6.2) is the same as the proof of (5.2), because $\sigma^{(1)}$ is the same for the Bochner and Hodge Laplacians.

Finally we compute $u_s^{(3)}$ using (5.18). From (6.4) and Lemma 4.4, we have

$$\begin{aligned} -\sigma^{(1)II} - 2\sigma^{(0)II} &= -(\xi \cdot h \cdot \xi) - |\xi|^2 \sum_i \operatorname{sgn}(i) h_{ii} - \frac{1}{2} \sum_{i,j} \operatorname{sgn}_I(i) \operatorname{sgn}_I(j) \xi_j^2 h_{ii} \\ -\sigma^{(1)II} + 2\sigma^{(0)II} &= -(\xi \cdot h \cdot \xi) + \frac{1}{2} \sum_{i,j} \operatorname{sgn}_I(i) \operatorname{sgn}_I(j) \xi_i^2 h_{jj} \\ (6.10) \quad \sum_j \sigma_{ij}^{(1)II} (\xi_i (h \cdot \xi)_j + \xi_j (h \cdot \xi)_i) &= 2|h \cdot \xi|^2 + (\xi \cdot h \cdot \xi) \sum_i \operatorname{sgn}_I(i) h_{ii}. \end{aligned}$$

and so putting this into (5.18), we get

$$\begin{aligned} \langle h, u_s^{(3)} h \rangle &= (2S-1)|\xi|^{n-2s-4} \sum_I \left((\xi \cdot h \cdot \xi) \left(-(\xi \cdot h \cdot \xi) - |\xi|^2 \sum_i \text{sgn}_I(i) h_{ii} - \frac{1}{2} \sum_{i,j} \text{sgn}_I(i) \text{sgn}_I(j) \xi_j^2 h_{ii} \right) \right. \\ &+ |\xi|^2 (\text{tr } h) \left(-(\xi \cdot h \cdot \xi) + \frac{1}{2} \sum_{i,j} \text{sgn}_I(i) \text{sgn}_I(j) \xi_j^2 h_{ii} \right) + |\xi|^2 \left(2|h \cdot \xi|^2 + (\xi \cdot h \cdot \xi) \sum_i \text{sgn}_I(i) h_{ii} \right) \Bigg). \end{aligned}$$

The two terms involving a single sign functions cancel, and we get

$$\begin{aligned} \langle h, u_s^{(3)} h \rangle &= (2S-1)|\xi|^{n-2s-4} \left(\binom{n}{p} (-(\xi \cdot h \cdot \xi)^2 - |\xi|^2 (\xi \cdot h \cdot \xi) (\text{tr } h) + 2|\xi|^2 |h \cdot \xi|^2) \right. \\ &\quad \left. + \frac{1}{2} (|\xi|^2 (\text{tr } h) - (\xi \cdot h \cdot \xi)) \sum_{i,j} \sum_I \text{sgn}_I(i) \text{sgn}_I(j) \xi_j^2 h_{ii} \right). \end{aligned}$$

From (7.2), this gives

$$\begin{aligned} \langle h, u_s^{(3)} h \rangle &= (2S-1)|\xi|^{n-2s-4} \left(\binom{n}{p} (-(\xi \cdot h \cdot \xi)^2 - |\xi|^2 (\xi \cdot h \cdot \xi) (\text{tr } h) + 2|\xi|^2 |h \cdot \xi|^2) \right. \\ &+ \frac{1}{2} \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) (|\xi|^2 (\text{tr } h) - (\xi \cdot h \cdot \xi)) |\xi|^2 (\text{tr } h) + 2 \binom{n-2}{p-1} (|\xi|^2 (\text{tr } h) - (\xi \cdot h \cdot \xi)) (\xi \cdot h \cdot \xi). \end{aligned}$$

Collecting terms, we get

$$\begin{aligned} \langle h, u_s^{(3)} h \rangle &= (2S-1)|\xi|^{n-2s-4} \left(2 \binom{n}{p} |\xi|^2 |h \cdot \xi|^2 + \left(-\binom{n}{p} - 2 \binom{n-2}{p-1} \right) (\xi \cdot h \cdot \xi)^2 \right. \\ &\quad \left. + \left(-\frac{3}{2} \binom{n}{p} + 4 \binom{n-2}{p-1} \right) |\xi|^2 (\text{tr } h) (\xi \cdot h \cdot \xi) + \frac{1}{2} \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) |\xi|^4 (\text{tr } h)^2 \right), \end{aligned}$$

which equals (6.3).

7. Free terms.

In Sections 5 and 6, we use the following elementary formulas. The sums here are over all subsets I of $\{1, \dots, n\}$ of size p .

$$(7.1) \quad \sum_I \chi_I(i) \chi_{I'}(j) = \binom{n-2}{p-1} (1 - \delta_{ij}).$$

$$(7.2) \quad \sum_I \text{sgn}_I(j) \text{sgn}_I(k) = \left(\binom{n}{p} - 4 \binom{n-2}{p-1} \right) + 4 \binom{n-2}{p-1} \delta_{jk}.$$

Proving such formulas is simple. For example, for (7.1), one considers separately the cases when $i = j$ and $i \neq j$. When $i \neq j$, the left hand side is the number of sets I which contain i but not j , which is $\binom{n-2}{p-1}$. When $i = j$ the left hand side is clearly zero. For (7.2), we first expand each sign function: $\text{sgn}_I = \chi_I - \chi_{I'}$, and then deal with each product of characteristic functions separately.

We end with a remark on another approach to carrying out the calculation of u_s . For the sums in (7.1) and (7.2), we refer to the coefficient on the right hand side which does not involve any delta functions as the *free term* of the sum. This is the value of the sum when all the fixed variables are distinct. For example, the free term of the sum on the left in (7.1) is $\binom{n-2}{p-1}$, and the free term of

$$(7.3) \quad \sum_I \text{sgn}_I(i) \text{sgn}_I(j) \text{sgn}_I(k) \text{sgn}_I(\ell)$$

is $16 \binom{n-4}{p-2} - 8 \binom{n-2}{p-1} + \binom{n}{p}$. If one tries to carry out the calculation of $\langle h, u_s h \rangle$ for the de Rham Laplacian without using coordinates in which h is diagonal, one is faced with several sums similar to (7.3). It turns out that one can replace such sums by their free terms, and the resulting answer is correct. The idea behind this is that the non-free terms always give rise to quadratic expressions in h which are not coordinate independent, whereas u_s is coordinate independent. If one could prove that the coordinate dependent terms arising from the non-free terms cannot add up in the end to produce a non-vanishing coordinate free term, then this would give another reasonably short way to carry out the calculation of u_s .

References

- [Ba] G. Baker and J. Dodziuk, Stability of spectra of Hodge-de Rham Laplacians, *Math. Z.* **224** (1997), no. 3, 327–345.
- [BGV] N. Berline, E. Getzler, and M. Vergne, *Heat kernels and Dirac operators*. Grundlehren der Mathematischen Wissenschaften, 298. Springer-Verlag, Berlin, 1992.
- [Br] T. Branson, Sharp inequalities, the functional determinant, and the complementary series. *Trans. Amer. Math. Soc.* **347** (1995), 3671–3742.
- [BCY] T. Branson, S. Y. A. Chang and P. Yang, Estimates and extremals for zeta function determinants on four-manifolds, *Commun. Math. Phys.* **149**, (1992), 241–262.
- [BØ] T. Branson and B. Ørsted, Conformal geometry and global invariants, *Differential Geometry and its Applications* **1**, 279–308, (1991).
- [BFKM] D. Burghelea, L. Friedlander, T. Kappeler and P. McDonald, On the functional logdet and related flows on the space of closed embedded curves on S^2 . *J. Funct. Anal.* **120**, (1994), 440–466.
- [CY] S. Y. A. Chang and P. Yang, Extremal metrics of zeta function determinants on 4-manifolds, *Annals of Math.* **142** (1995) 171–212.
- [CQ] S. Y. A. Chang and J. Qing, The zeta functional determinants on manifolds with boundary. *J. Funct. Anal.* **147** (1997), 327–399.

- [Ch] J. Cheeger, Analytic torsion and the heat equation, *Ann. Math.*, **109**, 259-322 (1979).
- [CT] S. Chaniillo, F. Treves, On the Lowest Eigenvalue of the Hodge Laplacian. *J. Diff. Geom.* (1997), **45**, 273-287.
- [CC] B. Colbois, G. Courtois, A Note on the First Nonzero Eigenvalue of the Laplacian Acting on p -forms, *Manuscripta Math.*, **68**, 143-160 (1990).
- [Do] J. Dodziuk, Eigenvalues of the Laplacian on Forms, *Proc. AMS*, **85**, 437-443 (1982).
- [Fe] H. Fegan, The spectrum of the Laplacian on forms over a Lie group. *Pacific J. Math.* **90** (1980), 373-387.
- [Go] R. Gornet, A New Construction of Isospectral Riemannian nilmanifolds with Examples, *Michigan Math J.* **43** (1996), no. 1, 159-188.
- [HZ] A. Hassell and S. Zelditch, Determinants of Laplacians in exterior domains. *Internat. Math. Res. Notices*, (1999), **18**, 971-1004.
- [Lo] J. Lott, Collapsing and the differential form Laplacian: the case of a smooth limit space. *Duke Math. J.* **114** (2002), 267-306.
- [Mo1] C. Morpurgo, Sharp trace inequalities for intertwining operators on S^n and R^n . *Internat. Math. Res. Notices* (1999), **20**, 1101-1117.
- [Mü] W. Müller, analytic torsion and R -torsion of Riemannian manifolds, *Adv. in Math.* **28** (1978), 233-305.
- [Ok1] K. Okikiolu, Critical metrics for the determinant of the Laplacian in odd dimensions. *Ann. of Math.* **153** (2001), 471-531.
- [Ok2] K. Okikiolu, Hessians of spectral zeta functions, *Duke Math. Journal*, to appear.
- [Ok3] K. Okikiolu, Critical metrics for the zeta functions of a family of scalar Laplacians, *Preprint*.
- [OPS1] B. Osgood, R. Phillips and P. Sarnak, Extremals of determinants of Laplacians, *J. Funct. Anal.* **80** (1988), 148-211.
- [OPS2] B. Osgood, R. Phillips, and P. Sarnak, Compact isospectral sets of surfaces. *J. Funct. Anal.* **80** (1988), 212-234.
- [OPS3] B. Osgood, R. Phillips, and P. Sarnak, Moduli space, heights and isospectral sets of plane domains. *Ann. of Math.* **129** (1989), 293-362.
- [Po] A. Polyakov, Quantum geometry of bosonic strings, *Phys. Lett. B* **103** (1981), 207-210.
- [RS] D. Ray and I. Singer, Analytic torsion, *Proc. Symposium Pure Math Vol. XXIII* AMS, Providence, R.I., 1973, 167-181.
- [Ri] K. Richardson, Critical points of the determinant of the Laplace operator, *Jour. Funct. Anal.*, **122**, (1994) 52-83.
- [Ro] S. Rosenberg, The variation of the de Rham zeta function, *Trans. AMS.* **299**, no. 2, (1987), 535-557.
- [Ta] J. Takahashi Upper bounds for the eigenvalues of the Laplacian on forms on certain Riemannian manifolds. *J. Math. Sci. Univ. Tokyo* **6** (1999), no. 1, 87-99.

Kate Okikiolu
University of California, San Diego
okikiolu@math.ucsd.edu

Caitlin Wang
University of California, San Diego
cwang@math.ucsd.edu